

RNA interference: An update on its application in insect control

Reshma, R., Sagar, D., and Rajna, S.

Abstract: RNA interference (RNAi) is being used as one of the important technologies for better designing of pest control strategies. It is found to be effective against different insect orders such as Coleoptera, Diptera, Lepidoptera and Hemiptera. The efficiency of RNAi depends on the design of dsRNA and the mode of delivery selected. The delivery of RNAi through transgenic plants is now a reality with some products currently in the market. Topically applied dsRNA/siRNA (Spray induced gene silencing) has attracted attention due to its feasibility & low cost compared to transgenic plants. Sustainable agriculture relies on practices and technologies that combine effectiveness with a minimal environmental footprint. RNAi-based products can revolutionize pest and pathogen management safely and effectively if conception and development are conducted in a precautionary way.

Key words: Gene silencing, Insects, delivery systems, dsRNA

An overview of RNAi in insects

RNAi is a post-transcriptional gene silencing mechanism initiated by introduction of double stranded RNA (dsRNA) leading to generation of loss of function phenotypes by degrading the target gene messenger RNAs (mRNAs). It utilises small RNA (sRNA) as trigger molecules for the manipulation of gene expression. sRNAs are derived from double stranded RNA (dsRNA) and come in a variety of forms each differing in structure, function and biogenesis. E.g, Micro RNA (miRNA, ~22nt long) which is transcribed from plant genomes to regulate expression of endogenous genes (Bosco et al, 2008) and small interfering RNA (siRNA, 20-25nt long) which is derived from exogenous dsRNA sequences or the products of miRNA directed silencing (Broughton *et al.*, 2014). The RNAi pathway involves cleavage of the introduced dsRNA by dicer within cells. The resulting short RNAs or siRNAs get unwound to guide strand and passenger strand, of which the latter degenerates. Guide strand gets incorporated to RNA-induced silencing complex (RISC). The RISC-RNA complex degrades the corresponding mRNA and reduce protein expression (Fig 1).

Sensitivity of insects to RNAi: The efficiency of RNAi varies with insects orders. Coleopterans show high sensitivity to dsRNA, Dipterans show moderate sensitivity while Lepidopterans and Hemipterans show weak sensitivity to dsRNA. This is due to the presence of dsRNA degrading enzymes in their digestive system. RNAi efficiency differs within insects of the same order and also based on the mode of delivery of dsRNA.

Uptake, release and export of dsRNA could affect RNAi efficiency and resistance in insects. Insects can uptake RNA as both dsRNA and siRNA, directly or indirectly. Direct uptake refers to uptake through topical contact or feeding on plant tissues while indirect uptake involves first entering RNA in to plant vascular system and then uptake by insects. The dsRNA reaches insect cells through SID-like transporters or endocytosis or with the help of specific receptors which are found more for long dsRNA when compared to siRNA. After entering the cell, dsRNA should escape from endosomes (internal sorting organelles) to get accessibility to target mRNA which is found to be limiting in some insects. Plants, fungi and nematodes have endogenous RNA dependent RNA polymerase (RdRp) which target

on single-stranded RNA molecules and synthesize a second strand, generating dsRNA consequently. This produces a systemic spread of RNAi signaling in them (Zotti *et al.*, 2017). Since RdRp is absent in insects further amplification of dsRNA is not possible. The dsRNA reaches from cell to cell (Systemic RNAi) through special structures like exosomes, nanotubes or other carrier molecules in insects (Karlikow *et al.*, 2016).

CHOICE OF dsRNA TARGETS IN INSECTS

The dsRNA targets selection can be of three types.

1. Resistance factors: Targeting genes associated with resistance to existing pesticides or other control measures. E.g, dsRNA targeting sodium channel increased sensitivity to pyrethroids in *Aedes aegypti* (Bona *et al.*, 2016), Asian citrus psyllid topically applied with dsRNA against *Cyo* genes increased sensitivity to imidacloprid (Kilny *et al.*, 2014).

2. Developmental and /or arthropod genes: Many insect-specific genes worth considering will reduce

the risk of off-target effects in vertebrates. E.g, Ecr KD in grain aphids (Yan *et al.*, 2016).

3. Intracellular trafficking pathways: These are tightly regulated processes used by a variety of molecules to cross the membranes of living cells. E.g, ESCART protein including *snf7* (Head *et al.*, 2017), *vATPases* (Baum *et al.*, 2007), COP pathway proteins (Taning *et al.*, 2018), Endocytosis proteins (Pinheiro *et al.*, 2018).

Besides these, dsRNA is also being studied for inducing sterility in insects (Whyard *et al.*, 2015).

DIFFERENT DELIVERY SYSTEMS OF dsRNA

After choosing the appropriate target genes, the most suitable delivery system has to be selected for the efficient application of RNAi in pest control. In laboratories, methods like artificial feeding and microinjection are utilized. For application of RNAi in fields, methods like Host-induced gene silencing (HIGS), Spray-induced gene silencing (SIGS), Virus-induced gene silencing (VIGS), Stem injection and Root absorption are used. Except for the first method, all others are non-transformative delivery systems.

1. Host-induced gene silencing (HIGS) (Transgenic in planta delivery): Entails the creation of transgenic crops that express the dsRNA specific for the pest.

SmartStax Pro is the first commercial RNAi product targeting on insect pests. It is a transgenic corn crop developed by Monsanto (Bayer crop science) against Western corn rootworm (WCR). It employs a pyramid strategy utilizing two different *Bt* proteins (Cry34Ab1/Cry35Ab1 and Cry3Bb1) as well as dsRNA targeting *snf7* gene expressed in the plant (Head *et al.*, 2017). *Bt* protein inserts itself into the gut epithelium and cause gut paralysis and thereby death. Down regulation of *snf7* gene, which plays an essential role in protein trafficking, will also bring mortality of the insect (Bolognesi *et al.*, 2012). This combined strategy lead to the swift death of the pest and less resistance development against the Plant Incorporated Protectant (PIP) (Head *et al.*, 2017). *Bt* genes also ensure protection from lepidopteran pests like fall armyworm resulting in a healthy crop both above and below ground. This product was approved by the United States Environmental Protection Agency in 2017. In 2016, it was released in Canada and a year later, in 2017, released in the USA.




Fig 1: Molecular mechanism of siRNA silencing. Source: <http://www.uni-konstanz.de/FuF/chemie/jhartig/>

for insects is more acceptable when compared to the latter.

Flock House Virus (FHV) was engineered to express *Drosophila melanogaster* specific dsRNA (Taning *et al.*, 2018). The advantage of using plant infecting virus is that it can move inside plant systematically through phloem, so recombinant virus can target phloem-feeding insects. Since VIGS is transiently transformed it does not cause alteration of the plant genome. Recombinant TMV expressing RNAi effectors infected *Nicotiana benthamina* plants caused death of citrus mealybug feeding on them (Khan *et al.*, 2013). This method is beneficial for woody plants since it takes time to produce GM crops. VIGS can be transmitted to plant progeny & actively co-opts the plant to express dsRNA. Virus simulate natural path of cell entry which is an additional benefit of this method. Cross infection of beneficial insects also challenges biosafety. Since viruses are not accepted widely due to these drawbacks, Virus-like particles (VLP) produced by certain microbes is an alternative.

VIGS-like technology is also employed with other microorganisms like bacteria, yeast, entomopathogenic fungi, microalgae etc. The use of microorganisms ensures sustained release of dsRNA. It also prevents rapid degradation of dsRNA in the digestive system. Since common bacteria like *Escherichia coli* are engineered and used, this bacteria may cause infection on beneficial insects too which is found to be one of the drawbacks. Thus, the use of symbiotic bacteria specific for particular insects is more preferred. Besides using bacteria as the delivery system, they are used to produce a large amount of dsRNA which can be sprayed on crops at low costs. The major drawback of VIGS is that it is considered as a GM product and goes through all related regulatory processes.

Comparison of delivery as naked dsRNA, through *E. coli* and symbiotic bacteria *Rhodococcus rhodnii* in kissing bug *Rhodnius prolixus* was successfully done by Whitten *et al.* (2017). They reported that all these methods invoke a systemic response. Though naked dsRNA and *E. coli* produced dsRNA resulted in only transient systemic RNAi (since *E. coli* get eliminated from the cell after dsRNA ingestion), symbiotic bacteria *Rhodococcus rhodnii* colonised the gut and thereby established sustained systemic RNAi. The engineered *R. rhodnii* were also transmitted horizontally through

ingestion of faeces which was found to be an additional benefit.

4. Stem injection and root absorption: Both are non-transgenic *in planta* delivery systems. Stem injection was found to be efficient for perennial trees. Hunter *et al.* (2012) reported that citrus trees treated with dsRNA through stem injection and root drench have shown effective control against citrus psyllid and leafhoppers for up to 57 days. Rice plant roots soaked in a solution containing dsRNA targeting *carboxylesterase* and *CYP18A1* genes from brown planthopper significantly knocked down these genes (Li *et al.*, 2015). Delivery of dsRNA molecules through irrigation water is an alternative for crops that use irrigation in normal growing systems (Li *et al.*, 2015). Though it allows a constant supply of dsRNA, the short persistence of dsRNA is a drawback associated with this method. Therefore, the use of advanced formulations is needed to protect dsRNA from degradation.

OTHER APPLICATIONS OF RNAi BASED PRODUCTS

Besides using against crop pests, RNAi based products are also developed for mosquito control, the human disease vector. A study on the efficiency of genetically engineered yeast expressing interfering RNA corresponding to mosquito neural genes as lure-and-kill mosquito and oviposition attractants were conducted and reported to be successful (Hapairai *et al.*, 2017). RNAi based products against pests of beneficial insects were also developed. Deformed Wing Virus infection on both larvae and adults were reduced by feeding European honeybees with dsRNA (Desai *et al.*, 2012). RNAi was also efficient against internal microsporidian parasite, *Nosema* (Paldi *et al.*, 2010) and obligatory ectoparasite *Varroa destructor* in *Apis mellifera* (Garbian *et al.*, 2012). Since dsRNA are target specific, they are found to be harmless against honeybees, so safety is assured.

RNAi BASED PRODUCTS: ACCOMPLISHMENTS

Currently approved RNAi based GM crops are based on ncRNA (non-coding RNA) to control insects (8%) and diseases (27%) or to improve scientific plant traits (65%). With all the drawbacks that GM products raise, more research is focusing on non-transformative delivery of dsRNA for gene silencing.

Monsanto is developing the use of RNAi through a technology called “Biodirect” in which dsRNA formulation is applied exogenously to protect plants against insect and pathogen attacks (<https://monsanto.com/innovations/agricultural-biologicals>). Syngenta is also developing lines of biocontrol products based on RNAi to protect potato plants from the attack of the colorado potato beetle (<https://www.youtube.com/embed/BiVZbAy4NHw?ecver=1>).

Since the production of inexpensive RNA is a primary need, a biotechnology start-up working on sprayable RNAi based insecticides *RNAissance Ag* introduced a technology in which industrial fermenting bacterial species are used to produce RNA in safe and cost-effective manner. Their sprayable RNAi based biopesticide against diamondback moth is under early field trial. Considering the hostile environmental condition to which dsRNA molecules are exposed in the field, the biotech company RNAgri (former APSE) developed a system where APSE RNA containers (ARC) are produced by *E. coli* bacteria, allowing the mass production of encapsulated ready to spray dsRNA. Research is being carried out on improving the potential of different RNAi based products and reducing their limitations by scientists from various countries.

During the last decade several researchers have carried work on RNAi based products, still there are some areas left behind. At field level, the time of application and rate of application of dsRNA has to be investigated. Since it is difficult to establish dsRNA uptake under field condition, new methods has to be discovered. The selection of most appropriate target gene, species specificity, cost of dsRNA, delivery of dsRNA formulations and regulation are major challenges while considering about RNAi based biopesticides. In India, since a more scientifically informed approach is followed for regulation of GM crops, topically applied RNAi based products will be a viable alternative. Recent study on topical application of dsRNA against papaya ring spot virus - Tirupati and Delhi isolate was successful providing 94% and 81% resistance respectively (Vadlamudi *et al.*, 2020). More studies in SIGS are in need against other economic pests also.

CONCLUSION

RNAi based products have immense potential in managing crop pests, pests which are a threat to beneficial insects, pathogens and insect vectors of human and plant diseases safely and effectively. Although GM products are not widely accepted, non-transformative approaches in RNAi can be exploited as pest control measures. Realization of the complete benefits of the technique can be achieved only through appropriate dsRNA design and delivery mechanisms. The major limitation associated with the products is their less stability and impact on non-target organisms. The use of suitable formulations will reduce the degradation of dsRNA in both environments and inside the host. Bioinformatics based design of dsRNA sequences to minimize homology with endogenous transcripts in both the host plant and NTOs is an important approach to mitigate the risk of biosafety (species-specific products). Synthesis of broad-spectrum biopesticides targeting a wide range of pests can also be explored in this way.

REFERENCES

- Baum J A, Bogaert T, Clinton W, Heck G R, Feldmann P, Ilagan O, Roberts J. 2007. Control of coleopteran insect pests through RNA interference. *Nature Biotechnology*. 25(11):1322-1326.
- Bernstein E, Caudy A A, Hammond S M, Hannon G J. 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature*. 409(6818):363-366.
- Bolognesi R, Ramaseshadri P, Anderson J, Bachman P, Clinton W, Flannagan R, Segers G. 2012. Characterizing the mechanism of action of double-stranded RNA activity against western corn rootworm (*Diabrotica virgifera virgifera* LeConte). *Plos one*, 7(10): e47534.
- Bosco L, Giacometto E, Tavella L. 2008. Colonization and predation of thrips (Thysanoptera: Thripidae) by *Orius* spp. (Heteroptera: Anthocoridae) in sweet pepper greenhouses in Northwest Italy. *Biological control*. 44(3):331-340.
- Broughton S, Harrison J, Rahman T. 2014. Effect of new and old pesticides on *Orius armatus* (Gross)—an Australian predator of western flower thrips, *Frankliniella occidentalis* (Pergande). *Pest*

- Management Science. 70(3):389-397.
- Cagliari D, Dias N P, Galdeano D M, dos Santos E A, Smaghe G, Zotti M J. 2019. Management of Pest Insects and Plant Diseases by Nontransformative RNAi. *Frontiers in Plant Science*, 10:1319.
- Christiaens O, Tardajos M G, Martinez Reyna Z L, Dash M, Dubruel P, Smaghe G. 2018. Increased RNAi efficacy in *Spodoptera exigua* via the formulation of dsRNA with guanylated polymers. *Frontiers in Physiology*. 9:316.
- Dalla Bona A C, Chitolina R F, Fermino M L, de Castro Poncio L, Weiss A, Lima J B P, Maori E. 2016. Larval application of sodium channel homologous dsRNA restores pyrethroid insecticide susceptibility in a resistant adult mosquito population. *Parasites & Vectors*. 9(1):1-14.
- Dalmay T, Hamilton A, Rudd S, Angell S, Baulcombe DC. 2000. An RNA-dependent RNA polymerase gene in *Arabidopsis* is required for posttranscriptional gene silencing mediated by a transgene but not by a virus. *Cell*. 101(5):543-53.
- Desai S D, Eu Y J, Whyard S, Currie R W. 2012. Reduction in deformed wing virus infection in larval and adult honey bees (*Apis mellifera* L.) by double stranded RNA ingestion. *Insect Molecular Biology*. 21(4):446-455.
- Fire A, Xu S, Montgomery M K, Kostas S A, Driver S E, Mello C C. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*. 391(6669):806-811.
- Fletcher S J, Reeves P T, Hoang B T, Mitter N. 2020. A perspective on RNAi-based biopesticides. *Frontiers in Plant Science*. 11:51.
- Gillet F X, Garcia R A, Macedo L L, Albuquerque E V, Silva M, Grossi-de-Sa M F. 2017. Investigating engineered ribonucleoprotein particles to improve oral RNAi delivery in crop insect pests. *Frontiers in Physiology*. 8:256.
- Hamilton A J, Baulcombe D C. 1999. A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science*: 286(5441):950-952.
- Hapairai L K, Mysore K, Chen Y, Harper E I, Scheel M P, Lesnik A M, Duman-Scheel M. 2017. Lure-and-kill yeast interfering RNA larvicides targeting neural genes in the human disease vector mosquito *Aedes aegypti*. *Scientific Reports*. 7(1):1-11.
- Head G P, Carroll M W, Evans S P, Rule D M, Willse A R, Clark T L, Meinke L J. 2017. Evaluation of SmartStax and SmartStax PRO maize against western corn rootworm and northern corn rootworm: efficacy and resistance management. *Pest Management Science*. 73(9):1883-1899.
- Hunter W B, Glick E, Paldi N, Bextine B R. 2012. Advances in RNA interference: dsRNA treatment in trees and grapevines for insect pest suppression. *Southwest Entomologist*. 37(1):85-87.
- Jacobsen S E, Running M P, Meyerowitz E M. 1999. Disruption of an RNA helicase/RNase III gene in *Arabidopsis* causes unregulated cell division in floral meristems. *Development*. 126(23):5231-5243.
- Karlikow, M., Goic, B., Mongelli, V., Salles, A., Schmitt, C., Bonne, I., Zurzolo, C. and Saleh, M.C., 2016. *Drosophila* cells use nanotube-like structures to transfer dsRNA and RNAi machinery between cells. *Scientific reports*, 6(1), pp.1-9.
- Khan, A.M., Ashfaq, M., Kiss, Z., Khan, A.A., Mansoor, S. and Falk, B.W. 2013. Use of recombinant tobacco mosaic virus to achieve RNA interference in plants against the citrus mealybug, *Planococcus citri* (Hemiptera: Pseudococcidae). *PLoS One*, 8(9), p.e73657.
- Killiny N, Kishk A. 2017. Delivery of dsRNA through topical feeding for RNA interference in the citrus sap piercing sucking hemipteran, *Diaphorina citri*. *Archives of Insect Biochemistry and Physiology*. 95(2):e21394.
- Li H, Guan R, Guo H, Miao X. 2015. New insights into an RNAi approach for plant defense against piercing sucking and stem borer insect pests. *Plant, Cell & Environment*. 38(11): 2277-2285.
- Liu J, Carmell M A, Rivas F V, Marsden C G, Thomson J M, Song J J, et al. 2004. Argonaute2 is the catalytic engine of mammalian RNAi. *Science*. 305:1437-1441.
- Mysore K, Flannery E M, Tomchaney M, Severson D W, Duman-Scheel M. 2013. Disruption of *Aedes aegypti* olfactory system development through chitosan/siRNA nanoparticle targeting

- of semaphorin-1a. PLoS neglected tropical diseases. 7(5):e2215.
- Petek M, Coll A, Ferenc R, Razinger J, Gruden K. 2020. Validating the potential of double-stranded RNA targeting Colorado potato beetle mesh gene in laboratory and field trials. *Frontiers in Plant Science*. 11:1250.
- Pinheiro, D.H., Velez, A.M., Fishilevich, E., Wang, H., Carneiro, N.P., Valencia-Jimenez, A., Valicente, F.H., Narva, K.E. and Siegfried, B.D. 2018. Clathrin-dependent endocytosis is associated with RNAi response in the western corn rootworm, *Diabrotica virgifera virgifera* LeConte. *PLoS one*, 13(8), p.e0201849.
- Taning, C.N., Christiaens, O., Li, X., Swevers, L., Casteels, H., Maes, M. and Smagghe, G. 2018. Engineered flock house virus for targeted gene suppression through RNAi in fruit flies (*Drosophila melanogaster*) in vitro and in vivo. *Frontiers in physiology*, 9, p.805.
- Vadlamudi T, Patil BL, Kaldis A, Gopal DV, Mishra R, Berbati M, Voloudakis A. 2020. DsRNA-mediated protection against two isolates of Papaya ringspot virus through topical application of dsRNA in papaya. *Journal of virological methods*. 275:113750.
- Vogel E, Santos D, Mingels L, Verdonck T W, Broeck J V. 2018. RNA Interference in Insects: Protecting beneficials and controlling pests. *Frontiers in Physiology*. 9:1912.
- Whyard S, Singh A D, Wong S. 2009. Ingested double-stranded RNAs can act as species-specific insecticides. *Insect Biochemistry and Molecular Biology*. 39(11):824-832.
- Whyard S, Erdelyan CN, Partridge AL, Singh AD, Beebe NW, Capina R. 2015. Silencing the buzz: a new approach to population suppression of mosquitoes by feeding larvae double-stranded RNAs. *Parasites & vectors*. 8(1):1-1.
- Wingard S A. 1928. Hosts and symptoms of ring spot, a virus disease of plants. *Journal of Agricultural Research*. 37:0127-0153.
- Xu P, Zhang Y., Kang L, Roossinck M J, and Mysore K S. 2006. Computational estimation and experimental verification of off-target silencing during posttranscriptional gene silencing in plants. *Plant physiology*. 142: 429–440.
- Yan T, Chen H, Sun Y, Yu X, Xia L. 2016. RNA interference of the ecdysone receptor genes EcR and USP in grain aphid (*Sitobion avenae* F.) affects its survival and fecundity upon feeding on wheat plants. *International Journal of Molecular Sciences*. 17(12):2098.
- Zhang X, Zhang J, Zhu K Y. 2010. Chitosan/double stranded RNA nanoparticle mediated RNA interference to silence chitin synthase genes through larval feeding in the African malaria mosquito (*Anopheles gambiae*). *Insect Molecular Biology*, 19(5):683-693.
- Zotti, M., Dos Santos, E.A., Cagliari, D., Christiaens, O., Taning, C.N.T. and Smagghe, G. 2018. RNA interference technology in crop protection against arthropod pests, pathogens and nematodes. *Pest management science*, 74(6):1239-1250.

AUTHORS

Reshma, R* (Corresponding author), Sagar, D. and Rajna, S., Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi 110 012

*Email: ramreshma11@gmail.com
