

GENERAL ARTICLE

Host Plant Associations in Butterflies: Ecological Importance and Conservation Significance

Diksha Chopra

Department of Zoology and Environmental Sciences, Punjabi University, Patiala-147002, Punjab, India.

chopradiksha31@gmail.com

Abstract

Butterflies are among the most conspicuous and widely appreciated insects, yet their survival depends on more than the presence of nectar-bearing flowers. The larval stages of butterflies require suitable host plants for feeding, growth and successful development. Host plant availability strongly influences oviposition and larval survival. Ultimately, it shapes the local distribution, population persistence, and overall diversity of butterflies. Many butterfly species are closely associated with one or a few plant taxa, while others use a wider range of hosts. These relationships shape not only the ecology of individual species but also broader patterns of community structure and habitat occupancy. In India, where both butterfly and plant diversity are high, host plant associations provide an important foundation for understanding butterfly biology and for developing effective conservation strategies. This article discusses the ecological role of larval host plants in butterflies, influence on diversity and distribution and their significance in habitat management and conservation.

Keywords: Butterflies, host plants, larval ecology, oviposition, diversity, conservation.

Introduction

Butterflies hold a special place in entomology because they are ecologically important, visually attractive and relatively easy to observe in the field. They are often used in ecological studies because their distribution and abundance reflect the quality and composition of surrounding vegetation. However, butterfly ecology cannot be understood fully by focusing only on adults. The butterfly life cycle includes egg, larva, pupa and adult stages and the successful completion of this cycle depends on the availability of appropriate resources at each stage (Dennis *et al.*, 2004; Tiple *et al.*, 2011).

Among these resources, larval host plants are of central importance. Female butterflies do not lay eggs randomly on vegetation. Instead, they

select plant species that can support larval feeding and development. This choice is crucial because newly hatched larvae are often highly dependent on the plant selected by the female and have limited ability to move in search of better food. Thus, host plant choice directly affects survival, growth and reproductive success (Dennis *et al.*, 2004).

Host plant relationships also influence broader ecological patterns. They affect local distribution, seasonal occurrence, habitat suitability and population stability. A site may contain abundant nectar plants and still fail to support breeding populations if larval host plants are absent. For this reason, host plant associations are fundamental not only to butterfly biology but also to conservation and habitat restoration (Tiple

et al., 2011; Nitin *et al.*, 2018).

Host Plants and Butterfly Life History

Host plants are the plants on which butterflies lay eggs and on which caterpillars feed after hatching. In many species, host use is highly specific. Some butterflies utilize only one plant species or a narrow range of closely related plants, whereas others can feed on a broader set of hosts. This variation in host specificity is one of the defining features of butterfly natural history (Dennis *et al.*, 2004). The process begins with oviposition. Female butterflies identify suitable plants using a combination of plant form, surface texture, colour, microhabitat and especially chemical cues. Plant secondary compounds often play a major role in host recognition. A plant that appears suitable to a human observer may be chemically unacceptable to a butterfly, whereas a less conspicuous native plant may be the preferred host. Such selectivity is adaptive because larval development depends heavily on plant suitability (Dennis *et al.*, 2004).




Figure 1: Adult butterfly visiting a flowering plant for nectar feeding in its natural habitat. (Photo: Diksha Chopra at Shimla, H.P.)

Host plants also determine larval performance. Nutritional value, water content, leaf toughness and defensive chemistry can influence feeding rate, growth, development time



Figure 2: Butterfly larva feeding on foliage, highlighting the importance of larval host plants in butterfly development. (Photo: Neeraj at HP)

and survival. Poor-quality host plants may delay development and increase larval exposure to predators and parasitoids. In contrast, suitable host plants support rapid growth and successful pupation. Therefore, host plant availability is not simply about food presence; it is about access to the correct food under the correct ecological conditions (Tiple *et al.*, 2011).

Host Specificity, Diversity and Butterfly Assemblages

Host plant associations shape butterfly diversity at several levels. At the species level, host specificity often determines habitat occupancy and geographic distribution. Specialist species are usually more vulnerable to habitat disturbance because they depend on a narrow set of plant resources. Generalist species may persist more easily in modified landscapes because they can use a wider range of plants. This difference partly explains why some butterfly species decline rapidly with habitat change while others remain common (Dennis *et al.*, 2004).

At the community level, host plant diversity contributes to butterfly richness. Comparative studies have shown a strong positive relationship between the diversity of host plants and the richness of butterfly species. Ferrer-Paris *et al.* (2013) demonstrated that butterfly species richness is strongly correlated with host plant diversity across higher taxonomic levels. This finding suggests that plant diversity provides the ecological foundation for butterfly diversity, since a wider variety of host plants can support a broader butterfly assemblage.

Host use also has evolutionary significance. Butterfly diversification has long been linked to plant associations and more recent network-based analysis suggest that both host shifts and the retention of ancestral host use capacity have contributed to butterfly diversification. Braga *et al.* (2018) showed that host-associated diversification is driven not only by shifts to new plants but also by variation in host use and butterfly plant network structure.

Evidence from Indian Studies

Indian studies illustrate the importance of host plant associations particularly well. In a detailed study from Nagpur city, Tiple *et al.* (2011) documented larval host plants for 120 butterfly species and recorded 126 host plants. Their work showed that butterflies in a tropical urban landscape depend on a wide range of plant forms, including trees, herbs, shrubs, climbers and parasitic plants. The study also demonstrated that host specificity, egg batch size and the number of host plants used were associated with butterfly population patterns in the city.

A broader perspective comes from the Western Ghats, one of the world's major biodiversity hotspots. Nitin *et al.* (2018) compiled an updated checklist showing that about 834 plant species or

species groups belonging to 88 families are used by 320 butterfly species in the Western Ghats. This checklist reveals the extraordinary scale and complexity of butterfly plant relationships in India and highlights the dependence of both endemic and widespread species on specific larval food plants. These studies show that host plant research is not merely descriptive. It provides essential insight into breeding biology, resource use, landscape ecology and conservation priorities. In the Indian context, where butterfly diversity varies across forests, grasslands, scrub, wetlands, agricultural mosaics and urban habitats, host plant documentation remains a highly valuable form of natural history research (Tiple *et al.*, 2011; Nitin *et al.*, 2018).

Field Observations on Host Plant Associations from Northwest India

Field observations from Northwest India further highlight the importance of larval host plants in determining butterfly occurrence and habitat use. During surveys, several lycaenid butterflies were recorded in association with their larval host plants and characteristic habitats. *Pratapa icetas* was recorded from Kareri Village, Himachal Pradesh, where its known larval host plant *Scurrula ferruginea* was present in the habitat. *Horaga onyx* was recorded from Himachal Pradesh and Uttarakhand, where its documented larval host plant *Coriaria nepalensis* occurs. The Common Silverline, *Cigaritis vulcanus*, was documented from different localities with reported host plants including *Ziziphus* spp. and *Plectronia parviflora*. Field observations also revealed habitat preferences linked with vegetation structure. *Loxura atymnus* showed a preference for shaded habitats near bamboo vegetation in the Doon Valley, whereas *Satyrium sassanides* was observed in high-elevation montane habitats of Lahaul Spiti. These observations emphasize that conservation

Table 1: Larval host plant associations of Indian butterflies.
Source: Nitin et al. (2018), and documented host plant records from literature.

Butterfly species	Common name	Butterfly family	Major larval host plant species (representative)	Host plant family
<i>Arhopala amantes</i>	Large Oak-blue	Lycaenidae	<i>Terminalia paniculata</i> , <i>Hopea ponga</i> , <i>Shorea robusta</i> , <i>Syzygium cumini</i>	Combretaceae, Dipterocarpaceae, Myrtaceae
<i>Danaus chrysippus</i>	Plain Tiger	Nymphalidae	<i>Calotropis gigantea</i> , <i>Calotropis procera</i>	Apocynaceae
<i>Delias eucharis</i>	Indian Jezebel	Pieridae	<i>Dendrophthoe falcata</i> , <i>Scurrula parasitica</i>	Loranthaceae
<i>Horaga onyx</i>	Common Onyx	Lycaenidae	<i>Mangifera indica</i>	Anacardiaceae
<i>Junonia lemonias</i>	Lemon Pansy	Nymphalidae	<i>Barleria cristata</i> , <i>Hygrophila auriculata</i>	Acanthaceae
<i>Pieris canidia</i>	Cabbage White	Pieridae	<i>Brassica oleracea</i> , <i>Rorippa indica</i>	Brassicaceae
<i>Rapala maene</i>	Slate Flash	Lycaenidae	<i>Mangifera indica</i> , <i>Acacia pennata</i> , <i>Saraca asoca</i>	Anacardiaceae, Fabaceae
<i>Surendra quercetorum</i>	Common Acacia Blue	Lycaenidae	<i>Acacia pennata</i> , <i>Albizia odoratissima</i>	Fabaceae

of butterfly diversity requires protection of both adult habitats and larval host plant resources.

Sorrel Sapphire, *Heliophorus sena*, has been recorded using *Rumex hastatus* as its larval host plant. Among the oakblues, the Sahyadri Rosy Oakblue, *Arhopala alea*, uses *Terminalia paniculata*, *Hopea* spp. and *Syzygium salicifolium*, whereas the Large Oakblue, *Arhopala amantes*, uses several tree species belonging to the families Combretaceae, Dipterocarpaceae, Fabaceae, Lythraceae and Myrtaceae. Common and widespread butterflies also show recognisable host associations: the Lime Swallowtail, *Papilio demoleus*, commonly uses citrus and other Rutaceae, while the Plain Tiger, *Danaus chrysippus*, breeds mainly on milkweeds such as *Calotropis* spp (Robinson et al., 2010, Kalesh and Prakash, 2015).

Conservation Significance

The conservation of butterflies is often approached by emphasizing nectar-rich flowering plants for adults. Although nectar plants are important, this approach is incomplete if it neglects larval host plants. A habitat may attract adult butterflies for feeding but still fail to support breeding populations if females cannot find appropriate host plants for oviposition. Effective conservation therefore requires a life-cycle approach in which both adult and larval resources are maintained (Tiple et al., 2011; Nitin et al., 2018). This issue is particularly important in modified landscapes. Urbanization, roadside clearing, replacement of native vegetation with ornamental plants and habitat simplification may reduce host plant availability even when green cover remains.



Figure 3: Butterfly larva associated with foliage, representing the dependence of immature stages on suitable plant resources. (Photo: Diksha Chopra at Shimla, H.P.)

Such habitats may appear suitable visually but may function poorly for butterfly reproduction. In many cases, the loss of larval host plants is one of the most important hidden causes of butterfly decline (Dennis *et al.*, 2004). Conservation planning should therefore prioritize native vegetation and locally relevant host plants. Restoration efforts in gardens, campuses, parks and reserve edges can be improved by including plant species known to support local butterfly larvae. This is especially important for specialist butterflies, which may depend on a limited set of host taxa. Because host plant diversity is linked to butterfly richness, protecting plant diversity is also a practical strategy for conserving butterfly communities more broadly (Ferrer-Paris *et al.*, 2013).

Natural History Value and Future Scope

Host plant associations remain one of the most rewarding areas of butterfly natural history. Careful observation of egg laying behaviour, larval feeding, resting sites, shelter-building and pupation can reveal valuable ecological information. Even simple field observations can contribute significantly when accompanied by accurate plant identification and photographic documentation.

There is also considerable scope for future work in India. Many regional host plant associations remain poorly documented and there is need for more field-based studies across seasons, elevations and habitat types. Changes in land use and climate may alter both plant availability and butterfly behaviour, making long-term observations increasingly important. Continued collaboration among professional researchers, amateur naturalists, students and citizen scientists could greatly improve available knowledge on butterfly host use in India (Nitin *et al.*, 2018).

Conclusion

Host plant associations are central to the ecology and conservation of butterflies. They influence oviposition, larval survival, development, habitat use and long-term population persistence. Evidence from both global comparative studies and Indian field research shows that butterfly diversity is closely tied to host plant diversity. For this reason, butterfly conservation should not focus solely on adult nectar resources. Equal attention must be given to the plants that sustain immature stages. A stronger host plant perspective will improve butterfly research, strengthen natural history documentation and lead to more effective conservation planning.

References

- Braga, M. P., Guimarães, P. R., Jr., Wheat, C. W., Nylin, S., & Janz, N. (2018). Unifying host associated diversification processes using butterfly-plant networks. *Nature Communications*, 9, 5155. <https://doi.org/10.1038/s41467-018-07677-x>
- Dennis, R. L. H., Hodgson, J. G., Grenyer, R., Shreeve, T. G., & Roy, D. B. (2004). Host plants and butterfly biology: Do host plant strategies drive butterfly status? *Ecological*

- Entomology, 29(1), 12-26. <https://doi.org/10.1111/j.1365-2311.2004.00572.x>
- Ferrer-Paris, J. R., Sánchez-Mercado, A., Vilorio, Á. L., & Donaldson, J. (2013). Congruence and diversity of butterfly host plant associations at higher taxonomic levels. PLoS ONE, 8(5), e63570. <https://doi.org/10.1371/journal.pone.0063570>
- Kalesh, S. & Prakash, S. K. (2015). Additions to the larval host plants of butterflies of the Western Ghats, Kerala, South India (Rhopalocera, Lepidoptera). Journal of Bombay Natural History Society, 112, 111-114.
- Nitin, R., Balakrishnan, V. C., Churi, P. V., Kalesh, S., Prakash, S., & Kunte, K. (2018). Larval host plants of the butterflies of the Western Ghats, India. Journal of Threatened Taxa, 10(4), 11495-11550. <https://doi.org/10.11609/jott.3104.10.4.11495-11550>
- Robinson, G. S., Ackery, P. R., Kitching, I. J., Beccaloni, G. W., & Hernández, L. M. (2010). HOSTS—a Database of the World's Lepidopteran Hostplants. Natural History Museum, London.
- Tiple, A. D., Khurad, A. M., & Dennis, R. L. H. (2011). Butterfly larval host plant use in a tropical urban context: Life history associations, herbivory, and landscape factors. Journal of Insect Science, 11(1), Article 65. <https://doi.org/10.1673/031.011.6501>