

GENERAL ARTICLE

The Silent Sculptor inside the Tobacco Stem - *Scrobipalpa aptatella* (Walker, 1864)

Guru, P. N.^{1*}, Raghavendra K. P.¹, Nanda, C.¹, Ramakrishnan, S.¹, Krithika, C. K.¹, Korada, R. R.¹ and Madhav, M. S.²

¹ICAR-National Institute for Research on Commercial Agriculture (NIRCA), R.S. Hunsur, Karnataka, India

²ICAR-National Institute for Research on Commercial Agriculture (NIRCA), Rajahmundry, Andhra Pradesh, India

*gurupn5016@gmail.com

Introduction

When farmers walk through a newly planted tobacco field in the early morning, the crop often appears healthy from a distance. The leaves sway gently in the breeze, the stems stand upright, and the field looks calm and green. But inside a few innocent-looking plants, a secret drama is unfolding. Deep within the stem lives a tiny hidden artist — the tobacco stem borer, *Scrobipalpa aptatella* (Walker, 1864). Unlike caterpillars that openly chew leaves or beetles that attack from outside, this insect prefers darkness. It lives silently inside the stem, tunneling, feeding, growing, and transforming without being seen.

The invisible thief

Like a silent thief inside a house, the stem borer eats the plant from within while the farmer notices nothing at first. The story usually begins when a tiny moth lays eggs on tender tobacco plants. Soon, the young larva hatches and searches for a safe entrance. Instead of feeding openly, it bores directly into the stem. Once inside, the insect becomes a hidden tenant. As days pass, the larva feeds internally on soft tissues. The plant reacts to this invasion by producing abnormal growth around the injured area. Slowly, the basal portion

of the stem swells and becomes enlarged like a knot or gall (Fig. 1a). To the farmer, this bulged region looks strange — almost as if the stem has developed a tumor. But inside that swollen chamber, the larva is comfortably feeding and preparing for its next stage of life (Jagadish, 1979). The gall is not just a symptom; it is the insect's secret house, which looks like a mysterious swelling at the base.

The stem first becomes food, shelter, nursery, and finally a coffin (not exactly as the adult emerges out from this secret house freely). The plant becomes its prison for two important stages i.e., larva and pupa.

An insect that fears fresh air

Most living creatures crave for air i.e. oxygen for survival. Humans need fresh air every moment. Animals struggle without oxygen. Even underground insects create breathing tunnels. But the tobacco stem borer is different. This insect survives only as long as it remains hidden inside the stem. If the swollen gall is carefully pierced or split open, exposing the larva to direct air and sunlight, the larva quickly dies (Fig. 1b). What an irony of nature! The same air that gives life to humans becomes deadly to this concealed insect. The very stem that protects the insect also becomes its trap




Figure 1. Various life stages and characteristic symptoms of damage caused by *Scrobipalpa aptatella* in tobacco (a) Gall at the basal portion of the stem (b) Larva inside the split open stem (c) Pupation site inside the stem (d) Multiple galls/ swellings on stem

when exposed to air. Farmers in some regions traditionally pin or puncture the swollen stem portion after noticing fresh galls. Once exposed, the delicate larva cannot tolerate the sudden environmental change. Nature itself becomes the pesticide. This simple observation teaches an important lesson: sometimes understanding an insect's weakness is more powerful than spraying chemicals. The stem borer survives in darkness, hidden from the world. But the moment sunlight and fresh air enter its chamber, its life ends.

The secret transformation inside the stem

As the larva matures, it prepares for one of nature's greatest miracles — metamorphosis. Inside the same stem tunnel, the caterpillar changes into a pupa. During this stage, it remains inactive, silently transforming into an adult moth. The pupation region often becomes more swollen and corky (Fig.1c). If the stem is cut open carefully, one can observe the brown pupa resting inside like a passenger in a sleeping chamber. Then comes the final act. One day, the adult moth emerges by cutting a neat circular hole through the stem wall. The hole looks as though someone drilled it using

a tiny machine (Rao and Rao, 1956; Sarala *et al.*, 2022). The moth escapes into the outside world, leaving behind an empty tunnel and a damaged plant. The adult moth cuts an almost perfect round hole as an escape door while emerging.

Does one insect attack only one plant?

A common question among farmers is: Does a single stem borer remain in one plant throughout its life? The answer is mostly yes during the larval stage. Once the larva enters a tobacco stem, it generally stays inside that single plant because the stem provides food, shelter, and protection. It completes feeding and pupation within the same stem (Fig. 1d). However, the adult moth emerges out from the gall is capable of laying eggs on many plants during the season. A single female moth can lay eggs on several tobacco plants. Thus, while one larva usually damages one plant, the next generation can spread widely across the field. However, if the multiple larvae emerges can able to bore the same plant where eggs laid, and we found up to 3 larvae (galls) in a single plant. That is why early detection is extremely important. One unnoticed gall today may become the source

of many infestations tomorrow. You can explain that one swelling alone can reveal feeding, growth, pupation, and adult emergence.


Why the damage is serious?

The stem is the backbone of the tobacco plant. When the insect tunnels inside; uptake gets disrupted, plant vigor declines, growth becomes stunted, leaves remain smaller, this may also cause horizontal growth or breakage near the basal portion of the plant (Sreenivas, 2022). In some cases, the damage can further expose the plant to secondary infection, particularly by *Fusarium* species, resulting in wilt disease. The gall formation also weakens the stem structurally.

In severe infestations, yield and leaf quality may reduce significantly. Since the pest hides internally, farmers often notice damage only after the swelling appears.

Nature's tiny engineer

Though harmful, the tobacco stem borer is also an example of nature's astonishing engineering. Imagine this tiny insect: entering a narrow stem, building tunnels, creating a protected chamber, transforming inside, and finally carving a perfect circular exit hole. All this happens silently inside a single plant stem. To a scientist, it is fascinating; To a farmer, it is alarming; and to nature, it is simply



survival.

A short life: A lasting impact

Interestingly, the life of the tobacco stem borer is also closely tied to its hidden lifestyle. The adult moth lives only for a short period, but during that time the female lays eggs on several tobacco plants across the field, helping the pest spread from one generation to another. Unlike some insects that reproduce only once in a dramatic final act, the moth may continue laying eggs over a few days before dying naturally. After mating and egg laying, the quiet cycle begins again — eggs hatch, larvae bore into fresh stems, new galls develop, and another hidden generation grows silently inside the plant.

Waiting silently between seasons – off-season survival

Even after the tobacco crop is harvested and the field appears empty, the story of the stem borer does not completely end. The insect quietly survives the off-season by hiding inside leftover stem pieces, stubbles, dried plant residues, and sometimes in alternate host plants growing nearby (Jagadish, 1979). Larvae or pupae concealed within these dry stems remain protected from harsh weather and enemies. When the next tobacco season begins, the adult moths emerge from these hidden shelters and start laying eggs on young plants once again. Thus, an unnoticed infested stem left in the field can become the starting point for a fresh infestation in the following season. This is why sanitation is extremely important in tobacco cultivation (Rao and Rao, 1956). Removal and destruction of old stalks, deep ploughing after harvest, and cleaning field borders help break the hidden bridge that carries the insect from one season to the next. In many ways, the stem borer spends the off-season like a patient sleeper inside dry stems, waiting quietly for the return of a new tobacco crop.

Nature also sends hunters

Although the tobacco stem borer hides safely inside the stem, it is not completely beyond nature's reach. Tiny parasitoid wasps, such as *Trichogramma* spp. (Rao and Sreedhar, 2013) and *Chelonus heliopae* (Patel *et al.*, 1958) attack the pest's eggs and early developmental stages, while predatory insects and wandering spiders constantly search tobacco fields for hidden prey. Some natural enemies attack the eggs or young larvae before they enter the stem, while others wait for the vulnerable stages during emergence. Thus, even this secret insect is part of a larger ecological battle taking place silently in the field. Nature always maintains balance — for every hidden pest, there is often a hidden hunter waiting nearby.

Why chemicals often fail

Farmers are often surprised when insecticide sprays do not give satisfactory control against tobacco stem borer. The reason lies in the insect's hidden lifestyle. By the time swelling becomes visible and spraying begins, the larva is already deep inside the stem, protected like a soldier hiding safely inside a bunker. The stem walls shield the insect from direct contact with chemicals, making late-stage control difficult. This is why early detection, field sanitation, and exposing the grub by splitting or pinning the swollen stem are often more effective than repeated spraying alone.

Outsmarting the hidden enemy – Management of tobacco stem borer

The tobacco stem borer teaches farmers an important lesson: a hidden enemy must be detected early. Since the insect spends most of its life safely inside the stem, ordinary spraying often fails once the larva enters the plant. Therefore, careful observation in the field becomes the farmer's greatest weapon.

- Clean cultivation also plays an important role. Removal of old crop residues, volunteer tobacco plants, and nearby weeds helps reduce carryover of the pest from one season to another.
- The moment a swollen gall-like bulge appears near the basal stem region, the plant should be examined closely. Experienced farmers sometimes gently pierce or split the swollen portion using a pin, thorn, or sharp stick.
- Plants showing severe swelling or drying should be removed and destroyed before the adult moth emerges. Otherwise, the moth escapes through its neat circular exit hole and flies away to lay eggs on many more tobacco plants across the field.
- Balanced fertilizer use should be encouraged because excessive nitrogen creates soft succulent growth that may attract more infestation.
- Farmers who regularly walk through their fields, observe stems carefully, and remove early infestations often succeed better than those depending only on chemicals. In many ways, management of tobacco stem borer is not merely spraying against an insect — it is understanding the secret life of a creature that lives hidden inside the stem.
- Under severely infested situations, after pinning the galls, the fields can be sprayed with recommended insecticides (Sreedhar, 2014).

Why scientists find the Tobacco Stem Borer fascinating

To a farmer, the tobacco stem borer is a damaging pest. But to scientists, it is also a remarkable example of hidden insect adaptation

and survival. Its entire life revolves around living inside a narrow plant stem — a tiny world where it feeds, grows, transforms, and finally emerges as an adult moth. Studying such an insect helps researchers understand how insects manipulate plant tissues, avoid predators, survive harsh environments, and complete their life cycle within a confined space. One of the most fascinating features is the gall-like swelling formed on the stem. Scientists are interested in how such a small larva can trigger the plant to alter its growth and create a protective chamber around it. The insect also demonstrates extraordinary behavioural adaptation: it survives safely in darkness inside the stem but becomes highly vulnerable when suddenly exposed to open air. The perfectly circular exit hole made by the adult moth is another marvel of natural engineering. Researchers study how the insect senses the correct direction for emergence and cuts through the stem wall at the right time without becoming trapped inside. The pest is also important in ecological and pest management studies because it remains hidden from many conventional insecticide sprays. Understanding its behavior helps scientists design better integrated pest management strategies using early detection, biological control, and cultural practices rather than depending only on chemicals. In many ways, the tobacco stem borer is not merely a crop pest — it is a living example of how insects evolve specialized lifestyles to survive inside plants.

Final reflection

The story of *Scrobipalpa aptatella* (Gelechiidae: Lepidoptera) reminds us that not all enemies attack openly. Some remain hidden, silently feeding from within. Yet this insect also carries a strange weakness: it survives in darkness but dies in open air. Perhaps nature is telling us something profound — every hidden problem

loses strength when exposed to light. And in every tobacco field, beneath the green leaves and quiet breeze, this small unseen drama continues season after season.

References

- Jagadish, A. (1979). Control of *Spodoptera litura* and *Scrobipalpa heliopa* in nursery tobacco. PANS (International Journal of Pest Management), 25(1), 16-18.
- Patel, H. K., Patel, R. C., & Patel, R. M. (1958). Some observations on *Chelonus heliopae* Gupta, a parasite of the tobacco stem borer. Indian Tobacco, 8(4), 233-235.
- Rao, S. G., & Sreedhar, U. (2013). Egg parasitoids in tobacco ecosystem. In *Biological Control of Insect Pests Using Egg Parasitoids* (pp. 317-329). New Delhi: Springer India.
- Rao, V. T., & Rao, N. R. (1956). The tobacco stemborer (*Phthorimoea heliopa* Low.) in the Northern Circars: and its control. Indian Journal of Entomology, 18(1), 49-56.
- Sarala, K., Nanda, C., Baghyalakshmi, K., Darvishzadeh, R., Prabhakara Rao, K., Sreedhar, U., & Gangadhara, K. (2022). Designing tobacco genomes for resistance to biotic stresses. In *Genomic Designing for Biotic Stress Resistant Technical Crops* (pp. 441-581). Cham: Springer International Publishing.
- Sreedhar, U. (2014). Management of tobacco stem borer, *Scrobipalpa heliopa* in virginia tobacco. Indian Journal of Plant Protection, 42(1), 6-10.
- Sreenivas, S. S. (2022). Influence of climatic change on pests and diseases of flue-cured Virginia (FCV) tobacco in India—Need for potential strategies. International Journal of Agriculture, Environment and Biotechnology, 15(3), 699-707.